

# DiffEqBiological Tutorial III: Steady-States and Bifurcations

Torkel Loman and Samuel Isaacson

June 29, 2020

Several types of steady state analysis can be performed for networks defined with DiffEqBiological by utilizing homotopy continuation. This allows for finding the steady states and bifurcations within a large class of systems. In this tutorial we'll go through several examples of using this functionality.

We start by loading the necessary packages:

```
using DiffEqBiological, Plots
gr(); default(fmt = :png);
```

## 0.0.1 Steady states and stability of a biochemical reaction network.

Bistable switches are well known biological motifs, characterised by the presence of two different stable steady states.

```
bistable_switch = @reaction_network begin
    d,      (X,Y) → ∅
    hillR(Y,v1,K1,n1), ∅ → X
    hillR(X,v2,K2,n2), ∅ → Y
end d v1 K1 n1 v2 K2 n2
d = 0.01;
v1 = 1.5; K1 = 30; n1 = 3;
v2 = 1.; K2 = 30; n2 = 3;
bistable_switch_p = [d, v1, K1, n1, v2, K2, n2];
```

```
7-element Array{Float64,1}:
 0.01
 1.5
30.0
 3.0
 1.0
30.0
 3.0
```

The steady states can be found using the `steady_states` function (which takes a reaction network and a set of parameter values as input). The stability of these steady states can be found using the `stability` function.

```
ss = steady_states(bistable_switch, bistable_switch_p)
```